

Infrared Colors at the Stellar/Substellar Boundary

S.K. Leggett

Joint Astronomy Centre, University Park, Hilo, HI 96720
skl@jach.hawaii.edu

F. Allard

CRAL, Ecole Normale Supérieure, 46 Allée d'Italie, Lyon, 69364 France
fallard@cral.ens-lyon.fr¹

and

P.H. Hauschildt

Department of Physics and Astronomy & Center for Simulation Physics,
University of Georgia, Athens, GA 30602-2451
yeti@hobbes.physast.uga.edu

full text with tables and figures available at
ftp://ftp.jach.hawaii.edu/pub/ukirt/skl/dM_preprint/

ABSTRACT

We present new infrared JHK photometry for 61 halo and disk stars around the stellar/substellar boundary. We also present new L' photometry for 21 of these stars and for 40 low-mass stars taken from the Leggett 1992 photometry compilation. These data are combined with available optical photometry and astrometric data to produce color-color and absolute magnitude-color diagrams — the current sample extends the similar work presented in the 1992 paper into more metal-poor and lower mass regimes. The disk and halo sequences are compared to the predictions of the latest model atmospheres and structural models.

We find good agreement between observation and theory except for known problems in the V and H passbands probably due to incomplete molecular data for TiO, metal hydrides and H₂O. The metal-poor M subdwarfs are well matched by the models as oxide opacity sources are less important in this case. The known extreme M subdwarfs have metallicities about one-hundredth solar, and the coolest subdwarfs have $T_{eff} \sim 3000$ K with masses $\sim 0.09 M_{\odot}$. The grainless models are not able

¹On leave from Wichita State University, Wichita, KS 67260

to reproduce the flux distributions of disk objects with $T_{eff} < 2500$ K, however a preliminary version of the NextGen–Dusty models which includes homogeneous formation and extinction by dust grains *is* able to match the colors of these very cool objects. The least luminous objects in this sample are GD165B, three DENIS objects — DBD0205, DBD1058 and DBD1228 — and Kelu-1. These have $T_{eff} \sim 2000$ K and are at or below the stellar limit with masses $\leq 0.075 M/M_{\odot}$. Photometry alone cannot constrain these parameters further as the age is unknown, but published lithium detections for two of these objects (Kelu-1 and DBD1228) imply that they are young (aged about 1 Gyr) and substellar (mass $\leq 0.06 M/M_{\odot}$).

1. Introduction

Low-mass stars are the most numerous type of object in the solar neighbourhood. Studies of the parameters of these long-lived objects — surface gravity, temperature, metallicity, luminosity, radius and mass — are important for our understanding of the Galactic mass density and its chemical evolution. In Leggett (1992, hereafter L92) photometric and kinematic data were compiled for about 300 red dwarfs and sequences were identified in color-color diagrams that were representative of the halo and the disk populations. These allowed empirical determinations to be made of metallicity and luminosity, based on observational colors.

Until very recently the observational data for low-mass stars could not be well reproduced by synthetic spectra or photometry. The cool, high-pressure atmospheres are difficult to model, due in particular to complex opacity sources: strong molecular bands and, for the halo stars and very low-mass objects, pressure-induced molecular hydrogen opacity (see e.g. Borysow, Jørgensen & Zheng 1997). The situation is now much improved as demonstrated by for example the “NextGen” models of Hauschildt, Allard & Baron 1998 (see also Allard et al. 1997). The observational side of the study of low-mass stars has also changed remarkably since 1992, with a large increase in the known number of low-mass stars, brown dwarfs, and even giant planets. This growth is due to the continuing dedicated effort by many groups, and to improvements in sensitivity and accuracy of the various techniques employed.

In this work we present new infrared photometry for a sample of halo and disk stars approaching and even below the stellar/sub-stellar boundary. This work extends L92 to more metal-poor and lower mass regimes. Also the data are taken with a well known photometric system, removing the need for uncertain color transformations. The disk and halo sequences are compared to synthetic colors from Allard & Hauschildt’s NextGen model atmospheres (Hauschildt, Allard & Baron 1998) and to isochrones from Baraffe & Chabrier’s structural models (Baraffe et al. 1997, Chabrier & Baraffe 1997) combined with the NextGen atmospheres. We also compare the data to colors from a preliminary version of NextGen which includes grain condensation (Allard et al. 1998a).

In §2 we describe the sample for which new photometry is presented in §3. §4 describes the model atmospheres and also describes the calculation of the synthetic photometry. In §5 the model predictions are compared to the observations, and metallicity, mass and effective temperature are estimated for the sample. Our conclusions are given in §6. The Appendix gives information on the CIT, IRTF and UKIRT photometric systems.

2. The Sample

JHK photometry was obtained for a selection of the more recently identified very low-mass stars (or brown dwarfs) of the halo and disk. The halo stars were selected from studies of known high proper-motion stars by Gizis 1997, Gizis & Reid 1997 and Monet et al. 1992. The disk stars were selected from publications of new discoveries by various color or proper-motion surveys — Delfosse et al. 1997 (the DENIS survey DBD objects); Irwin, McMahon & Reid 1991 (the BRI objects); Kirkpatrick, Beichman & Skrutskie 1997 (the 2MASS survey); Ruiz, Leggett & Allard 1997 (Kelu-1); Schneider et al. 1991 (PC0025); Tinney 1993 (the TVLM objects) — or from follow-up studies of known high proper-motion stars — Kirkpatrick, Henry & Irwin 1997; Monet et al. 1992 — or from studies of low-mass companions to nearby stars — Becklin & Zuckerman 1988 (GD165B). The sample also includes four red dwarf companions to cool white dwarfs observed in the course of the white dwarf work by Bergeron, Ruiz & Leggett 1997 (LP356-87, LHS 229, G111-72 and G59-39). The target stars are listed in Table 1. We give LHS or LP number (Luyten 1979), and/or Gliese or Gliese/Jahreiss number (Gliese & Jahreiss 1991), and/or Giclas number (Giclas, Burnham & Thomas 1971), for each star. We also give discovery names for the objects taken from the sources cited above, where we have abbreviated “TVLM” to “T”. An abbreviated RA/Dec is also given to aid identification.

The spectral types in Table 1 are taken from various sources and there may be discrepancies or errors at the level of one sub-class. For the halo stars the classifications primarily are from Gizis 1997 and Gizis & Reid 1997, but for LHS 2045, LHS 2245, LHS 3168 and LHS 3390 they are based on our own optical spectra. For the disk stars the classifications are taken from Gizis 1997, Kirkpatrick, Henry & Simons 1995 and Kirkpatrick, Henry & Irwin 1997, and we have classified the new objects Kelu-1 and the DENIS objects by this scheme, based on published optical spectra.

The sources for the astrometric data — trigonometric parallax and radial velocity — are given in Table 1. We have calculated the UVW space motions for each object that has both parallax and radial velocity measurements, and classified these objects into kinematic populations of young disk (YD), young/old disk (Y/O), old disk (OD), old disk/halo (O/H) and halo (H). We have followed the velocity specifications described in L92 for these classifications.

3. Observations

We obtained JHK data for the stars in Table 1 using infrared cameras on two telescopes, both situated on Mauna Kea in Hawaii. These cameras were NSFCAM on the NASA Infrared Telescope Facility (Shure et al. 1994) and IRCAM on the UK Infrared Telescope (Puxley et al. 1994) both of which use InSb detectors. The JHK L' filter sets for the instruments are effectively identical except for the H filter. The only color transformation required is at H and the correction onto the UKIRT system for those objects observed only with NSFCAM is $\leq 7\%$. We give the NSFCAM–IRCAM (IRTF–UKIRT) H–band color transformation in the Appendix. UKIRT Faint Standards were used to calibrate the JHK photometry and UKIRT Bright Standards were used for L' (Casali & Hawarden 1992, see also URL <http://www.jach.hawaii.edu/UKIRT/astronomy/standards.html>); the Appendix gives some more information on the UKIRT system.

Table 2 lists new JHK data for the 61 stars in our sample where we have converted the NSFCAM H–band data to the UKIRT system. We include data for LHS 377 although these data were obtained in 1994 and published in Leggett et al. 1996; we list it here again for reference, as it was not included in L92. The other data were obtained on various telescope runs over the period 1995 to 1998; the camera used and the number of observations are given in Table 2. The technique used was the standard “jitter” technique, also known as “dithering”. This technique involves making multiple exposures, typically 5, with the telescope offset between each exposure such that the star is placed on different regions of the array. Based on the reproducibility of the photometry between jitters the errors at JHK are 3% for stars with $K < 14$ and 5% for the fainter stars.

Table 2 also gives new L' photometry for 21 stars, and more new L' data are given in Table 3 for 40 stars in the L92 sample. These data were also taken with either NSFCAM or IRCAM, for which the filter and photometric systems are identical. The data were obtained during the period 1995 to 1998, except for LHS 377 and LHS 248 for which L' photometry was obtained through the UKIRT Service Observing Program in 1994. The L' data for these two stars were published in Leggett et al. 1996 but we include the data here for reference. Errors are given for L' for each star in Tables 2 and 3.

Table 4 gives VIJHK L' colors for the sample of 61 stars specified in Table 1. The optical data are taken from various publications as listed in the table, and are on the Cousins system (the data from Weis 1987 and Eggen 1979 were converted to the Cousins system using the relationships summarised in L92). We also give the distance modulus $M - m$ determined from the parallax given in Table 1.

4. Models

For comparison with the observed photometry we show the results of two sets of models, the grainless NextGen models (Hauschildt, Allard & Baron 1998) and the NextGen–Dusty models

(Allard et al. 1998a). These models have been computed with the stellar atmosphere code PHOENIX (version 9.1) in its static, plane-parallel, LTE mode (see e.g. Hauschildt, Baron & Allard 1997). The metallicities used are all simple scaled solar metallicity values, with, for example, no enhancement of α elements.

4.1. NextGen

The NextGen models are described in Allard et al. 1997 and Hauschildt, Allard & Baron 1998. The equation of state for these models is an enlarged and enhanced version of that presented in Allard & Hauschildt 1995. Also, new molecular line data have been incorporated. These include the lines from the HITRAN92 (Rothman et al. 1992) database, line lists for YO (Littleton 1987) and ZrO (Littleton & Davis 1985), the Jørgensen 1994 list of TiO lines (as well as other lists from his SCAN database), and the available list of water lines of Miller & Tennyson as described in Allard et al. 1994.

Both atomic and molecular lines are treated with a direct opacity sampling method. We do *not* use pre-computed opacity sampling tables, but instead dynamically select the relevant LTE background lines from master line lists at the beginning of each iteration for every model and sum the contribution of every line within a search window to compute the total line opacity at *arbitrary* wavelength points. In the calculations we present in this paper, we have included a constant statistical velocity field, $\xi = 2$ km/s, which is treated as a microturbulence. The choice of lines is dictated by whether they are stronger than a threshold $\Gamma \equiv \chi_l/\kappa_c = 10^{-4}$, where χ_l is the extinction coefficient of the line at the line center and κ_c is the local bound-free absorption coefficient. This typically leads to about $20 \cdot 10^6$ lines which are selected from the master line lists (with 47 million atomic and up to 340 million molecular lines). The profiles of these lines are assumed to be depth-dependent Voigt profiles (or Doppler for very weak lines). Details of the computation of the damping constants and the line profiles are given in Schweitzer et al. 1996.

4.2. NextGen-Dusty

The NextGen-Dusty models have been introduced in Allard et al. 1997 and a detailed publication is in progress (Allard et al. 1998a). These models include both the formation of and the extinction by dust grains (absorption and scattering) — they are present both in the equation of state and as an opacity source. The chemical equilibrium has been augmented to include, besides a complete gas phase composition of over 600 species, the formation of over a thousand crystal and liquid condensates. These so-called dust grains are assumed to be in equilibrium with the surrounding gas phase medium, and their abundance is computed using the Gibbs energy of formation compiled in the JANAF thermochemical tables (Chase et al. 1985).

These models assume an interstellar grain size distribution and reconstruct the grain opacity

coefficients as a function of wavelength using polarizability spectra from the literature and the Continuous Distribution of Ellipsoids theory of Bohren & Hoffman 1983. The opacity profiles (both absorption and scattering) of twelve types of grain have been included — Fe, Ni, Cu, corundum (Al_2O_3), MgAl_2O_3 , enstatite (MgSiO_3), forsterite (Mg_2SiO_4), and the silicates CaSiO_3 , Ca_2SiO_4 , $\text{CaMgSi}_2\text{O}_6$, $\text{Ca}_2\text{Al}_2\text{SiO}_7$, and $\text{Ca}_2\text{MgSi}_2\text{O}_7$. These are the most abundant condensates in late-type M dwarfs. The opacity of three of these grain species (Fe, forsterite, and $\text{Ca}_2\text{MgSi}_2\text{O}_7$) had however to be omitted to reproduce the infrared spectral distribution of late-type red dwarfs (see e.g. Ruiz, Leggett & Allard 1997), suggesting that these grains have settled gravitationally below the photosphere, or have failed the nucleation process. Further discussion is beyond the scope of this paper, but will be presented in Allard et al. 1998a. The Ruiz et al. work showed that inclusion of these particular species (and only these species) leads to an excessive amount of infrared flux compared to the observed energy distribution of Kelu-1 (the effect of grains is to heat the upper layers of the atmosphere and to disassociate H_2O). This work shows that Kelu-1 does not have unusual colors and these species must be omitted for all late-type red dwarfs.

A further improvement on the NextGen grid is the use of the latest water vapor line list by Partridge & Schwenke 1997 which includes over 300 million transitions, and is more complete for the higher energy transitions than the Miller & Tennyson line list. The Partridge & Schwenke list has been used for the models with $T_{eff} < 2000$ K and has the effect of increasing the depth of the water bands, compared to the less complete Miller & Tennyson line list (but this effect is offset by that of the inclusion of grains).

The NextGen–Dusty model results presented in this work are preliminary, but they do demonstrate the likely effect of the inclusion of grains in the atmospheres of very low mass stars and brown dwarfs. The diagrams in the following sections show that these preliminary dusty models do a very good job of reproducing the observed colors of low-mass objects, and that the grainless models fail for $T_{eff} < 2500$ K. As the dusty models are preliminary, we adopt single values of surface gravity and metallicity given by $\log g = 5.0$ and $[\text{m}/\text{H}] = 0$. Grains will also be important for lower metallicities; initial calculations show that for $[\text{m}/\text{H}] = -0.5$ condensation affects the spectral energy distribution for $T_{eff} < 2300$ K and for $[\text{m}/\text{H}] = -1.0$ it is important for $T_{eff} < 2000$ K. However there are no metal-poor stars in our sample at such low temperatures and we defer presentation of these results to Allard et al. 1998a.

4.3. Synthetic Photometry

The JHKL' colors on the UKIRT system were calculated by integrating the synthetic spectra over the UKIRT bandpasses (see the Appendix for more information on the UKIRT system). The VI color computations used the Cousins V and I bandpasses as given by Bessell 1990. Terrestrial atmospheric absorption was included in all filter bandpasses. The magnitudes were calibrated by dividing the synthetic spectra by the observed spectrum of Vega (Hayes 1985, Mountain et al. 1985), which was adopted to define zero magnitudes at all wavelengths.

4.4. Structural Models and Absolute Magnitudes

Structural and evolutionary models for very low mass stars have recently been calculated — by Baraffe et al. 1997 and Chabrier & Baraffe 1997 — which use updated physics for these dense and cool objects together with the NextGen (grainless) atmospheres. These isochrones provide a theoretical relationship between effective temperature and surface gravity, at a given metallicity, mass and age. Synthetic color grids have been calculated from the NextGen synthetic spectra for a range of effective temperature, metallicity and surface gravity, and these grids were interpolated to produce colors and absolute magnitudes at the appropriate effective temperature and gravity for a mass sequence at a given metallicity and age. Synthetic color grids have also been calculated from the NextGen–Dusty models but these have not yet been combined with the structural models, and for these models we adopt single values of surface gravity and metallicity given by $\log g = 5.0$ and $[m/H]=0$. Where absolute magnitudes are presented for these dusty models in the diagrams below, these magnitudes have been derived by scaling the calculated stellar surface magnitudes to match the NextGen (grainless) absolute magnitudes at 2900K. At this temperature condensation is not important (but below this temperature it is).

We have investigated whether gravity can be reliably determined from the observed photometry, as this could provide an independent check on the theoretical T_{eff} – $\log g$ relationship used in the evolutionary models, or a check on the age of the stars in the sample. For the disk stars the structural models imply that $\log g$ ranges from 4.7 at the hot/massive end to 5.4 at the cool/low-mass end, for an age of 10 Gyr. For a younger age of 1 Gyr, and solar metallicity, the low-mass value is slightly lower — $\log g \sim 5.3$. For the halo stars the range is $5.1 \leq \log g \leq 5.5$. We found that for a reasonably large change in gravity of 0.5 dex the synthetic colors differ by an amount less than the observational error, and so we cannot constrain the theoretical T_{eff} – $\log g$ relationship or the age of the stars.

5. Comparison of Data and Models

5.1. Color–Color Diagrams

In all the diagrams in this subsection we show colors generated by interpolating the NextGen model color grid onto the T_{eff} – $\log g$ values generated by model isochrones, as described above. We use a fixed age of 10 Gyr as colors alone are insensitive to varying age (while keeping metallicity constant), except for the lowest mass objects at very young ages (results for an age of 1 Gyr are shown in the absolute magnitude–color diagrams presented in the following section). The NextGen–Dusty colors have been generated for a single value of $\log g = 5.0$. This simplification to a single value for surface gravity is valid as the colors are not a strong function of gravity (for example, at $T_{eff} = 1900$ K changing $\log g$ by 0.5 dex only changes the various colors plotted in the following diagrams by ≤ 0.10 mag).

In all the plots the filled symbols are new data from the current work and the open symbols are from L92. The L92 data have been converted to the UKIRT JHK photometric system using transformations given by Casali & Hawarden 1992 (see Appendix). Symbol shapes illustrate kinematic populations: squares are halo, triangles are disk (including all young disk to old disk/halo) and circles are unknown kinematic population.

Figure 1 shows the V–I:I–K diagram with NextGen model sequences for metallicities $[m/H] = -2, -1$ and 0 and the NextGen–Dusty colors for $\log g = 5.0$ and $[m/H] = 0$. Lines of constant effective temperature are superposed. A typical observational error bar is shown, as well as the larger error bars for Kelu-1. It can be seen that while the metal-poor halo sequence appears to be well reproduced (the extreme subdwarf LHS 205a is labelled), there is a problem for the disk sequence. Other color–color diagrams show that the problem is in the V band, and in this spectral region there are known inadequacies in the metal hydride and titanium oxide spectroscopic data (Alvarez & Plez 1997). It is not known whether these molecular data are the source of the error, whether there is some other form of missing opacity, or whether the problem is structural, such as the treatment of convection. Although problems in one spectral region imply that an overall redistribution of the stellar flux is needed, very little flux is emitted at V for the red dwarfs — tests on the model atmospheres indicate that even a 0.5 mag additional opacity in the V bandpass would not cause significant flux redistribution to JHK wavelengths (Baraffe et al. 1998).

Figure 2 shows I–J:J–K which is a more complete representation of our data sample (only eight stars do not have I data available compared to twenty lacking V, some of which are the reddest objects). Again a typical error bar is shown as well as the larger errors for Kelu-1 and the DENIS objects. The cool extreme subdwarfs are labelled as well as the coolest disk objects. Note that the famous non-controversial brown dwarf Gl 229B (Nakajima et al. 1995) is not shown as it has very different colors due to strong methane absorption in the H and K bands ($I \sim 20.0$, $J \sim H \sim K \sim 14.3$, Matthews et al. 1996). Spectral types for the disk stars are given on the right-hand axis of Figure 5. These are based on the relationship between I–J and spectral class demonstrated in L92, and they agree with the individual classifications listed in Table 1. The recently discovered extremely red dwarfs have raised the issue of classification beyond M9, as discussed by Kirkpatrick 1998.

Although it is known that the Miller & Tennyson line list for water, the most important near-infrared opacity source, is incomplete (Allard et al. 1997), the agreement between theory and observation is good for these colors — which cover the peak in the stellar energy distribution. For the coolest objects the agreement becomes remarkably good when the effect of grain formation and extinction is taken into account. At the blue end of the sequence there is a discrepancy at the 10% level which is not seen if the L92 data is matched in CIT–system space via other filter profiles (Allard et al. 1998b). Either there is an error at this level in the models, which is cancelled by errors in the CIT color transformation, or there are problems with the UKIRT defined system. One potential source of error is the UKIRT J filter which has its red transmission edge determined by telluric water vapor absorption. However computations using a realistic range of telluric water

vapor show that photometric deviations of only about 3% are likely. This problem should be resolved with improvements to infrared photometric systems (such as the filter set now being purchased by a Mauna Kea Observatory consortium which is better matched to the atmospheric windows) and further refinements to the models.

Figure 3 is the J–H:H–K diagram in the UKIRT system. We also indicate the CIT system values on opposite axes for comparison to other published work. We have labelled the bluest (in J–H) metal-poor M dwarfs as well as the reddest disk objects. This diagram is still very difficult for the models to reproduce, at least for disk stars. The problem here is predominantly the H band where the overall opacity is lower, and hence photons are originating in a regime of higher temperature and pressure — where the Miller & Tennyson H₂O opacity profile is poorly defined. Nevertheless the models present a clear improvement in this diagram compared to previous model generations. Metal-poor models match the behavior of the most extreme subdwarfs, with colors becoming bluer as the effective temperature decreases (due to pressure-induced H₂ absorption in the K bandpass), while metal-rich models and stars are confined to red J-H values. Dust formation accounts for the extreme redness of the least massive metal-rich stars.

Figure 4 shows the longer wavelength J–K:K–L′. The outliers LHS 375 — an extreme subdwarf — and Kelu-1 — an extremely cool disk object — are labelled. L′ photometry is only available for the brighter members of this and the L92 sample. The agreement is good especially when grain opacities are included for the coolest objects.

5.2. HR Diagrams

In these diagrams the symbols are the same as in the earlier figures. In addition, large crosses show the location of the end of the main sequence, based on Baraffe et al. 1997 and Chabrier & Baraffe 1997 models. The coolest subdwarfs and disk objects are labelled.

The parallaxes for the L92 data (open symbols) have been updated using the final version of the Yale Catalogue (van Altena, Lee, & Hoffleit 1994) resulting in small changes to the observed absolute magnitudes for this sample. Vertical error bars are shown for stars with parallax errors greater than 20%. Note that these diagrams show that the parallax for the distant pair TVLM514-42404A/B has been overestimated. Only 50 of the 61 stars in the current sample with parallax measurements available are plotted, but not all of these appear on all the diagrams due to incomplete color coverage — the redder colors are more complete.

In absolute magnitude–color plots such as these the metal-poor stars form what appears to be a “sub-luminous” sequence sitting below the main sequence — hence the name “subdwarfs”. In fact, as is well known, the metal-poor halo stars are hotter and bluer than more metal-rich stars of the same mass, due to decreased optical and increased infrared opacity. The exact shift with metallicity is a function of the color plotted and the mass of the star. Although there is not a one-to-one correspondence between space motion and metallicity, the halo kinematic sample

clearly defines a “sub-luminous” sequence in these figures.

Figure 5 shows $M_V:V-I$ with 10 Gyr isochrones superimposed for metallicities $[m/H] = -2, -1$ and 0. A dotted line also shows the solar metallicity 1 Gyr isochrone, however this is indistinguishable from its older counterpart as no structural differences are expected except for masses lower than $0.09M/M_\odot$ and $M_V > 17$, at which point younger objects of the same mass will be brighter. Lines of constant mass are shown. The model discrepancy in the V band for the more metal-rich stars is again apparent.

Figure 6 shows the better reproduced $M_I:I-J$, and Figure 7 shows $M_J:J-K$ where the right axis shows empirical masses as a function of absolute J magnitude, based on the relationship determined by Henry & McCarthy 1993 for low-mass binary systems. These masses are valid for the disk stars only. The dotted line is the model sequence for an age of 1 Gyr with the hydrogen burning limit corresponding to brighter magnitudes compared to the older sequence. Despite the ~ 0.1 mag offset seen in J–K, as discussed in §5.1, the match between observation and theory is reasonable, especially when grains are included for the cool disk objects. The theoretical masses agree well with those derived observationally by Henry & McCarthy.

5.3. Results

Table 5 gives estimates of metallicity, mass and effective temperature for the stars in our sample, based on the model comparisons shown in Figures 1–7. These classifications were performed by eye, taking into account observational error and apparent systematic discrepancies between theory and observation at V–I and J–K, to produce a consistent estimate of metallicity and effective temperature. We have grouped the stars first into broad metallicity classes of $[m/H] \sim -2, -1$, and 0, and listed them in order of decreasing effective temperature within these groups. We indicate values of T_{eff} every 500 K, with the corresponding mass as implied by the structural models of Baraffe et al. 1997 and Chabrier & Baraffe 1997.

These values of metallicity, mass and effective temperature can only be estimates as spectroscopy is required for a more accurate determination of stellar parameters, and for the lowest mass objects age is also important (and unknown). Also, it is likely that the absolute values determined by model fitting will change in the near future as the models are upgraded. Nevertheless the overall trends indicated in Table 5 will be correct.

Spectroscopic lithium detections have been recently published for three of the coolest objects in this sample — Kelu-1 (Ruiz, Leggett & Allard 1997, Basri et al. 1997), DBD1228 (Basri et al. 1997, Tinney, Delfosse & Forveille 1997), and LP944-20 (Tinney 1998). The detection of lithium combined with determinations of $T_{eff} \sim 2000$ K for these three objects implies that their ages are about 1 Gyr and necessarily constrains their masses to be $\leq 0.06M/M_\odot$, i.e. substellar, based on current models by Chabrier & Baraffe 1997. This mass constraint arises from the fact that lithium depletion timescales are short, so for a star to show lithium it has to be either well below

the stellar mass limit or it has to be extremely young (aged about 0.1 Gyr). However if it is extremely young *and* massive enough to be stellar then Chabrier & Baraffe 1997 calculate that $T_{eff} > 2800$ K, i.e. much hotter than these objects. The age constraint arises from the fact that a substellar object would have cooled below 2000 K if the age were greater than about 1 Gyr. Finally, the fact that LP944-20 and Kelu-1 have partially depleted their lithium implies they are not significantly younger than 1 Gyr.

Basri et al. 1997 also looked for lithium in DBD1058 but failed to detect it. The depletion of lithium and the fact that $T_{eff} \sim 2000$ K, implies a mass for this object between $0.06M/M_{\odot}$, if it is aged about 1 Gyr, and $0.075M/M_{\odot}$ (the stellar limit), if it is aged several Gyr.

Amongst the metal-poor stars LHS 453 and LP 251-35 are notable as they are the most metal-poor stars in our sample with $[m/H] < -2$, based on the comparisons with the models presented in this work. These stars are known extreme subdwarfs. Another metal-poor star of particular interest is LHS 254, which is very red. Our photometry implies that it may be as metal-poor as LHS 377 with $[m/H] = -1$ and hence could be very close to the hydrogen-burning limit for that metallicity. Follow up observations are required, including measurement of its trigonometric parallax (such work is in progress at the US Naval Observatory Flagstaff Station).

6. Conclusions

The last two years have seen remarkable advances in the study of very low-mass stars and brown dwarfs. These advances have been on both the observational front, where we now have well defined sequences for both the halo and disk extending close to or beyond the brown dwarf boundary, and on the theoretical front, with much more complex and realistic models for both the stellar atmosphere and structure.

We have presented new infrared data for a sample of the low mass metal-poor stars and the very cool disk objects. These data are all on the same photometric system and allow the halo and disk sequences to be identified down to the hydrogen burning boundary. Although halo stars at the stellar limit have not yet been identified, the current sample includes halo red dwarfs with masses down to $\sim 0.09M/M_{\odot}$ and $T_{eff} \leq 3000$ K. The disk sample includes lithium-confirmed field brown dwarfs — Kelu-1 (Ruiz, Leggett & Allard 1997, Basri et al. 1997), DBD1228 (Basri et al. 1997, Tinney, Delfosse & Forveille 1997) and LP 944-20 (Tinney 1998).

Comparison to theoretical models gives good agreement in color-color diagrams, especially when grain condensation is included. Remaining problem areas are being worked on — improvements to the line lists for molecular species such as TiO, hydrides and H₂O, and inclusion of the gravitational settling of grains — and we can look forward to improvements in the match to the observational data in the near future. A more detailed spectroscopic comparison between observation and theory will be presented by this group in a subsequent paper.

We can also no doubt look forward to a continuing increase in the number of known low mass objects as observational efforts continue, including the large-scale sky surveys by the DENIS group (Delfosse et al. 1997) and 2MASS (Kirkpatrick, Beichman & Skrutskie 1997), which are now in full production mode.

We are very grateful to the staff at IRTF and UKIRT for their assistance in obtaining the data presented in this paper. UKIRT, the United Kingdom Infrared Telescope, is operated by the Joint Astronomy Centre on behalf of the U.K. Particle Physics and Astronomy Research Council. The IRTF, the NASA Infrared Telescope Facility, is operated by the University of Hawaii under contract to NASA. The UKIRT Service program also contributed to this work. Isabelle Baraffe kindly interpolated the atmospheric models on to the interior models to determine mass and radius at various ages, and Alastair Glasse calculated the transmission of the Mauna Kea atmosphere as a function of precipitable water vapor. We are grateful to Chris Tinney and Neill Reid for early access to their radial velocity data, and to Allyn Smith and Terry Oswalt for permission to use unpublished optical data. Tom Geballe read through a draft of the paper and gave useful comments. FA acknowledges support from NASA LTSA NAG5-3435 and NASA EPSCoR grants to Wichita State University. PHH acknowledges partial support by NASA ATP grant NAG 5-3018 and LTSA grant NAG 5-3619 to the University of Georgia. Some of the calculations presented in this paper were performed on the IBM SP2 of the UGA UCNS, at the San Diego Supercomputer Center (SDSC), the Cornell Theory Center (CTC), and at the National Center for Supercomputing Applications (NCSA), with support from the National Science Foundation. We thank all these institutions for a generous allocation of computer time.

A. Appendix — Infrared Color Transformations

In L92 all JHKL data were given on the CIT photometric system. At that time the CIT system was commonly used, with many infrared telescopes using the standard stars on this system published by Elias et al. 1982. Since 1992, however, more sensitive infrared array cameras have become the instruments of choice and the commonly used set of standards is now the UKIRT Faint Standards List published by Casali & Hawarden 1992 (see also URL <http://www.jach.hawaii.edu/UKIRT/astronomy/standards.html>). For this reason, as well as to avoid introducing errors through uncertain color transformations, we have left the new data presented in this work on the UKIRT system. Many color transformations were summarised in L92 and here we wish to update the CIT to UKIRT transformations. The most recent determination is that published by Casali & Hawarden 1992 and the following should be used in preference to the relations given in L92:

$$K_{CIT} = K_{UKIRT} - [0.018 \times (J - K)]$$

$$(J - K)_{CIT} = 0.936 \times (J - K)_{UKIRT}$$

$$(H - K)_{CIT} = 0.960 \times (H - K)_{UKIRT}$$

$$(J - H)_{CIT} = 0.920 \times (J - H)_{UKIRT}$$

Note that the original UKIRT Faint Standards data, and the transformation to CIT, were obtained using a single aperture photometer UKT9. However comparisons between UKT9 and IRCAM show no significant systematic difference for stars with $J - K < 1$ (Casali & Hawarden 1992; Hawarden, private communication).

Copies of the UKIRT filter profiles can be retrieved by following the “available filters” link from URL <http://www.jach.hawaii.edu/UKIRT/instruments/ircam/ircam3.html>. During the course of this work we determined that the IRTF’s NSFCAM filters are effectively identical to UKIRT’s IRCAM filters at J, K and L’, but not at H. Observations of red dwarfs show that the transformation required to bring IRTF–H onto the UKIRT system is:

$$(H - K)_{UKIRT} = 0.82[\pm 0.02] \times (H - K)_{IRTF}$$

which is valid for $0 \leq (H - K)_{UKIRT} \leq 0.5$.

REFERENCES

- Allard, F., 1998, “Model Atmospheres of Very Low Mass Stars and Brown Dwarfs”, in *The Tenth Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun*, (ed) A. Dupree, PASP Conference Series, in press
- Allard, F. & Hauschildt, P.H., 1995, ApJ, 445, 433
- Allard, F., Alexander, D. R., Tamanai, A., and Hauschildt, P. H., 1998b, “Photospheric Dust Grains Formation in Brown Dwarfs”, in *Brown Dwarfs and Extrasolar Planets*, (ed) R. Rebolo, PASP Conference Series, in press
- Allard, F., Alexander, D. R., Tamanai, A. & Hauschildt, P.H., 1998a, in preparation for ApJ
- Allard, F., Hauschildt, P.H., Alexander, D.R., & Starrfield, S., 1997, ARA&A, 35, 137
- Allard, F., Hauschildt, P.H., Baraffe, I. & Chabrier, G. 1996, ApJ, 465, L123
- Allard, F., Hauschildt, P. H., Miller, S., & Tennyson, J. 1994, ApJ, 426, 39
- Alvarez, R. and Plez, B., 1998, A&A, 330, 1109
- Baraffe, I., Chabrier, G., Allard, F. & Hauschildt, P. H., 1997, A&A, 327, 1054
- Baraffe, I., Chabrier, G., Allard, F. & Hauschildt, P. H., 1998, A&A, in press
- Basri, G., Martín, E., Ruiz, M.T., Delfosse, X., Forveille, T., Epchtein, N., Allard, F. & Leggett, S.K., 1998, “The First Look at Line Profiles in Cool Field Brown Dwarfs”, in *The Tenth Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun*, (ed) A. Dupree, PASP Conference Series, in press
- Becklin, E. E. and Zuckerman, B., 1988, Nature, 336, 656
- Bergeron, P., Ruiz, M.T., & Leggett, S.K. 1997, ApJS, 108, 339
- Bessell, M.S., 1990, PASP, 102, 1181
- Bessell, M.S., 1991, AJ, 101, 662
- Bohren, C.F. & Huffman, D.R., 1983, *Absorption and Scattering of Light by small Particles*, New York: Wiley & Sons, 541
- Borysow, A., Jørgensen, U.G. & Zheng, C., 1997, A&A, 324, 185
- Bouvier, L., Stauffer, J., Martin, E.L., Barrado y Navasues, D., Wallace, B., and Bejar, V. 1998, A&A, submitted
- Casali, M. & Hawarden T.G. 1992, UKIRT Newsletter Number 4, 33

- Chabrier, G. and Baraffe, I., 1997, A&A, 327, 1039
- Chase, M.W., Jr., Davies, C. A., Downey, J. R., Jr., Frurip, P. J., McDonald, R. A., Syverud, A. N., 1985, JANAF Thermochemical Tables, 3rd ed., J. Phys. Chem. Ref. Data 14, Suppl. No. 1, Parts 1 and 2
- Dawson, P.C. & de Robertis, M.M., 1988, AJ, 95, 1251
- Dawson, P.C. & de Robertis, M.M., 1989, AJ, 958 1472
- Dawson, P.C. & Forbes, D., 1989, PASP, 101, 614
- Delfosse, X. et al., 1997, A&A, 327, L25
- Eggen, O.J., 1979, ApJS, 39, 89
- Elias, J.H., Frogel, J.A., Matthews, K., & Neugebauer, G., 1982 AJ, 87, 1029
- Giampapa, M.S. & Liebert, J., 1986, ApJ, 305, 784
- Giclas, H.L., Burnham, R. & Thomas, N.G., 1971, Lowell Proper Motion Survey, The G Numbered Stars (Flagstaff: Lowell Observatory)
- Gizis, J.E., 1997, AJ, 113, 806
- Gizis, J.E. & Reid, I.N., 1997, PASP, 109, 849
- Gliese, W., & Jahreiss, H., 1991, Preliminary Version of the Third Catalog of Nearby Stars, Astronomisches Rechen-Institut Heidelberg Germany
- Harrington, R.S. & Dahn, C.C., 1980, AJ, 85, 454
- Hauschildt, P.H., Allard, F., & Baron, E., 1998, ApJ, submitted
- Hauschildt, P.H., Baron, E. & Allard, F., 1997, ApJ, 483, 390
- Hayes, D.S. 1985, in IAU Symp. 111, Calibration of Fundamental Stellar Quantities, ed. D.S. Hayes, E. Pasinetti & A.G.D. Philip (Dordrecht: Reidel), 225
- Henry, T.J. & McCarthy Jr., D.W., 1993, AJ, 106, 773
- Irwin, M., McMahon, R.G. & Reid, I.N., 1991, MNRAS, 252, 61p
- Jørgensen, U. G. 1994, A&A, 284, 179
- Kirkpatrick, J. D., 1998, “Brown Dwarfs and Extra-Solar Planets”, in *Brown Dwarfs and Extrasolar Planets*, (ed) R. Rebolo, PASP Conference Series, in press
- Kirkpatrick, J. D., Beichman, C.A. & Skrutskie, M.F., 1997, ApJ, 476, 311

- Kirkpatrick, J. D., Henry, T. J., & Irwin, M.J., 1997, *AJ*, 113, 1421
- Kirkpatrick, J. D., Henry, T. J., & Simons, D. A., 1995, *AJ*, 109, 797
- Leggett, S.K., 1992, *ApJS*, 82, 351
- Leggett, S.K., Allard, F., Berriman, G., Dahn, C.C. & Hauschildt, P.H., 1996, *ApJS*, 104, 117
- Leinert, CH., Weitzel, N., Richichi, A., Eckart, A., Tacconi–Garman, L. E. , 1994, *A&A*, 291, L47
- Littleton, J. E. 1987, private communication.
- Littleton, J. E., & Davis, S. P., 1985, *ApJ*, 296, 152
- Luyten W.J. 1979, *The LHS Catalogue, Second Edition* (Minneapolis: University of Minnesota)
- Matthews, K., Nakajima, T., Kulkarni, S.R., Oppenheimer, B.R., 1996, *AJ*, 112, 1678
- Monet, et al. 1992, *AJ*, 103, 638
- Mountain, C.M., Leggett, S.K., Selby, M.J., Blackwell, D.E. & Petford, A.D., 1985, *A&A*, 150, 281
- Nakajima, T., Oppenheimer, B.R., Kulkarni, S.R., Golimowski, D.A., Matthews, K. & Durrance, S.T. 1995, *Nature*, 378, 463
- Partridge, H., & Schwenke, D. W. 1997, *J. Chem. Phys.*, 106, 4618
- Puxley, P., Sylvester, J., Pickup, A., Paterson, M., Laird, D. & Atad, E., 1994, *Instrumentation in Astronomy VIII*, (eds Crawford, D.L & Craine, E.R.), *proc. SPIE* 2198, 350)
- Reid, I.N., Hawley, S.L., & Gizis, J.E., 1995, *AJ*, 110, 1838
- Rothman, L. S., et al., 1992, *JQSRT*, 48, 469
- Ruiz, M.T. & Anguita, C., 1993, *AJ*, 105, 614
- Ruiz, M.T., Leggett, S.K. & Allard, F., 1997, *ApJ*, 491, L107
- Schneider, D.P., Greenstein, J.L., Schmidt, M. & Gunn, J.E., 1991, *AJ*, 102, 1180
- Schweitzer, A., Hauschildt, P. H., Allard, F., & Basri, G. 1996, *MNRAS*, 283, 821
- Shure, M.A, Toomey, D.W, Rayner, J.T., Onaka, P. & Denault, A.J., 1994, *Instrumentation in Astronomy VIII*, (eds Crawford, D.L & Craine, E.R.), *proc. SPIE* 2198, 614
- Tinney, C.G., 1993, *ApJ*, 414, 279
- Tinney, C.G., Reid, I.N., Gizis, J.E. & Mould, J.R., 1995, *AJ*, 110, 3014
- Tinney, C.G., 1996, *MNRAS*, 281, 644

- Tinney, C.G., Delfosse, X. & Forveille, T., 1997, *ApJ*, 490, L95
- Tinney, C.G., 1998, *MNRAS*, 296, L42
- Tinney, C.G., & Reid, I.N., 1998, *MNRAS*, submitted (preprint available from <http://xxx.lanl.gov/ps/astro-ph/9806004>)
- van Altena, W.F., Lee, J.T., & Hoffleit, E.D. 1994, *The General Catalogue of Trigonometric Parallaxes* (New Haven: Yale University Observatory)
- Viti, S., Jones, H.R.A., Schweitzer, A., Allard, F., Hauschildt, P.H., Tennyson, J., Miller, S, Longmore, A.J., 1997, *MNRAS*, 291, 780
- Weis E.W. 1987, *AJ*, 93, 451

Fig. 1.— V–I:I–K diagram with model sequences for metallicities $[m/H] = -2, -1$ and 0 . Lines of constant effective temperature are shown. Filled symbols — this work, open — L92 on the UKIRT JHK system. Symbol shapes are kinematic populations: squares — halo, triangles — disk, circles — unknown.

Fig. 2.— I–J:J–K with model sequences using the same symbols as in Figure 1. Spectral types based on I–J (L92) are shown.

Fig. 3.— J–H:H–K diagram with model sequences using the same symbols as in Figure 1.

Fig. 4.— J–K:K–L' with model sequences using the same symbols as in Figure 1.

Fig. 5.— M_V :V–I with 10 Gyr isochrones superimposed for metallicities $[m/H] = -2, -1$ and 0 . Lines of constant mass are shown. The symbols are as in Figure 1. The dotted line is the 1 Gyr isochrone, and large crosses indicate the end of the hydrogen burning sequences.

Fig. 6.— M_I :I–J, with symbols and line types as in Figure 5.

Fig. 7.— M_J :J–K, with symbols and line types as in Figure 5. Empirical masses for the disk stars from Henry & McCarthy 1993 are shown.